

## Schriftenverzeichnis

1. Scherer R, Giebler R, **Kampe S**, Kox W. Effects of hypertonic saline hydroxyethyl starch solution on collagen-induced platelet aggregation and ATP secretion. *Transfusions Med Hemoth* 1994;21:310-14.
2. **Kampe S**, Weigand C, Kaufmann J, Klimek M, König DP, Lynch J. Postoperative analgesia with no motor block by continuous epidural infusion of ropivacaine 0.1% and sufentanil after total hip replacement. *Anesth Analg* 1999;89:395-8.
3. **Kampe S**, Iffland R, Korenkov M, Diefenbach C. Survival from a lethal blood concentration of cyanide with associated alcohol intoxication. *Anaesthesia* 2000;55:1-3.
4. **Kampe S**, Randebrock G, Kiencke P, Hünseler U, Cranfield K, König DP, Diefenbach C. Comparison of continuous epidural infusion of ropivacaine and sufentanil with intravenous patient-controlled analgesia after total hip replacement. *Anaesthesia* 2001;56:1189-93.
5. Kasper SM, Strömich A, **Kampe S**, Radbruch L. Evaluation of a new hydroxyethyl starch solution (HES 130/0.4) in patients undergoing preoperative autologous blood donation. *J Clin Anesth* 2001;13:486-90.
6. Krombach J, **Kampe S**, Gathof BS, Diefenbach C, Kasper SM. Human error: the persisting risk of blood transfusion: a report of five cases. *Anesth Analg* 2002;94:154-6.
7. **Kampe S**, Diefenbach C, Kanis B, Auweiler M, Kiencke P, Cranfield K. Epidural combination of ropivacaine with sufentanil for postoperative analgesia after total knee replacement. A pilot study. *Eur J Anaesthesiol* 2002; 19(9):666-71.
8. **Kampe S**, Kiencke P, Krombach J, Cranfield K, Kasper SM, Diefenbach C. Current practice in postoperative epidural analgesia: a German survey. *Anesth Analg* 2002;95:1767-9.
9. **Kampe S**, Nori H, Kasper SM, Diefenbach C. Unzureichende Ausbreitung einer Spinalanästhesie mit Bupivacain 0.5%/5% Glucose hyperbar im Vergleich zu einem Präparat von Bupivacain 0.5%/8% Glucose hyperbar. *Anaesthesist* 2002;51:993-5.

10. **Kampe S**, Nori H, Schneider P, Krott R. Unklare Mydriasis während einer Ösophagektomie mit unauffälligem Narkoseverlauf - Medikamentenwirkung von Akrinor? Ein Fallbericht. AINS 2003;38:165-7.
11. Kasper SM, Meinert P, **Kampe S**, Görg C, Geisen C, Mehlhorn U, Diefenbach C. Large-dose hydroxyethyl starch 130/0.4 does not increase blood loss and transfusion requirements in coronary artery bypass surgery compared with hydroxyethyl starch 200/0.5 at recommended doses. Anesthesiology 2003;99:42-7.
12. **Kampe S**, Poetter C, Buzello S, Wenchel HM, Paul M, Kiencke P, Kasper SM. Ropivacaine 0.1% with sufentanil 1 µg/mL inhibits growth of Pseudomonas aeruginosa and does not promote multiplication of Staphylococcus aureus. Anesth Analg 2003;97:409-11.
13. **Kampe S**, Kiencke P, Delis A, Auweiler M, König DP, Kasper SM. The continuous epidural infusion of ropivacaine 0.1% with 0.5 µg/mL sufentanil provides effective postoperative analgesia after total hip replacement: a pilot study. Can J Anaesth 2003;50:580-5.
14. Paul M, Dueck M, **Kampe S**, Petzke F. Failure to detect an unusual obstruction in a reinforced endotracheal tube with fiberoptic examination. Anesth Analg 2003;97:909-10.
15. **Kampe S**, Auweiler M, Kiencke P, Kasper SM, Diefenbach C. Wandel in der geburtshilflichen Periduralanalgesie? Eine bundesweite Umfrage aus dem Jahr 2000. Anästhesiol Intensivmed 2003;44:619-22.
16. **Kampe S**, Warm M, Kasper SM, Diefenbach C. Concept for postoperative analgesia after TRAM flap surgery: Continuous wound instillation with 0.2% ropivacaine via multilumen catheters. A report of two patients. Br J Plastic Surg 2003;56:478-83.
17. Paul M, Dueck M, **Kampe S**, Petzke F, Ladra A. Intracranial placement of a nasotracheal tube following transnasal transsphenoidal surgery. Br J Anaesth 2003 2003;91(4):601-4.
18. Paul M, Dueck M, **Kampe S**, Freundt H, Kasper SM. Pharmacological characteristics and side effects of a new galenic formulation of propofol without soybean oil. Anaesthesia 2003;58:1056-62.
19. **Kampe S**, Tausch B, Paul M, Kasper SM, Bauer K, Diefenbach C, Kiencke P. Comparison of epidural ropivacaine 0.75% and bupivacaine 0.5% for cesarean section: Cardiovascular stability and neonatal outcome. CMRO 2004;20(1):7-12.

20. Hekmat K, Zimmermann T, **Kampe S**, Weber HJ, Lauterbach G, Mehlhorn U. Impact of tranexamic acid vs aprotinin on blood loss and transfusion requirements after cardiopulmonary bypass: a prospective, randomised, double-blind trial. *CMRO* 2004;20(1):121-6.
21. **Kampe S**, Warm M, Kaufmann J, Hundegger S, Mellinghoff H, Kiencke P. Controlled-release oxycodone 20 mg administered on a 12-h dosing schedule halves postoperative IV opioid PCA consumption after breast surgery for cancer. *CMRO* 2004;20(2):199-202.
22. Kaufmann J, Yesiloglu S, Patermann B, Krombach J, Kiencke P, **Kampe S**. Controlled-release oxycodone is better tolerated than intravenous tramadol/metamizol for postoperative analgesia after retinal-surgery. *Curr Eye Res* 2004;28(4):271-5.
23. Patermann B, Buzello S, Dueck M, Paul M, **Kampe S**. Accidental tenfold overdose in a 6-month old infant undergoing craniosynostosis repair. *Anaesthesia* 2004;59(9):912-14.
24. Pilgram B, Krombach J, Ladra A, **Kampe S**. Relaxiert wach und trotzdem "relaxed" – ein Fallbericht. *Anästhesist* 2004;53(7):629-32.
25. Auweiler M, **Kampe S**, Zähringer, Buzello S, von Spiegel T, Buzello W. The human error: Delayed diagnosis of intravascular loss of guide wires for central venous catheterization. *J Clin Anesth* 2005;17(7):562-4.
26. Hekmat K, Kroener A, Schwinger RHG, **Kampe S**, Fischer UM, Geissler HJ, Mehlhorn U. The effect of preoperative antiplatelet therapy in coronary artery surgery: blood transfusion requirements for patients on cardiopulmonary bypass. *CMRO* 2004;20(9):1429-35.
27. Kaufmann J, Klimek M, **Kampe S**, Paul M, Kasper SM. A survey of autologous blood transfusion practices in Germany, 2000. *Transfusion Medicine* 2004;31:1-7.
28. Hekmat K, Kroener A, Stuetzer H, Schwinger RHG, **Kampe S**, Bennink GBWE, Mehlhorn U. Daily assessment of organ dysfunction and survival in ICU cardiac surgical patients. *Annals Thor Surg* 2005;79(5):1555-62.
29. Hekmat K, Kröner A, Stützer H, Schwinger RHG, **Kampe S**, Bennink GBWE, Mehlhorn. Ein neuer Score für die tägliche Schweregradklassifizierung auf herzchirurgischen Intensivstationen. *Zeitschrift für Herz- Thorax- und Gefäßchirurgie* 2004;18:257-67.
30. Landwehr S, Kiencke P, Giesecke T, Eggert D, Thumann G, **Kampe S**. A comparison between IV paracetamol and IV metamizol for postoperative analgesia after retinal surgery. *CMRO* 2005;21(10):1569-75.

31. Hekmat K, Raabe A, Kroener A, Fischer U, Suedkamp M, Geissler HJ, Schwinger RH, **Kampe S**, Mehlhorn U. Risk stratification models fail to predict hospital costs of cardiac surgery patients. *Z Kardiol* 2005;94(11):748-53.
32. Patermann B, Kiencke P, Fürderer S, König DP, Paul M, **Kampe S**. Patienten nach Spinalanästhesien haben einen höheren Quality of Recovery Score als Patienten nach Intubationsnarkosen – Klinische Anwendung des Quality of Recovery Scores. *Anästhesiol Intensivmed* 2007;2:64-70.
33. Schroeder S, von Spiegel T, Stueber F, Hoeft A, Preusse CJ, Welz A, **Kampe S**, Lier H. Interleukin 6 Enhancement after direct autologous retransfusion of shed thoracic blood does not influence hemodynamic stability following coronary artery bypass grafting. *Thoracic Cardiovasc Surg* 2007;55:68-72.
34. **Kampe S**, Warm M, Landwehr S, Dagtekin O, Haussamnn S, Paul M, Pilgram B, Kiencke P. IV paracetamol 1 g is clinically equivalent to IV dipyron for postoperative analgesia after breast surgery for cancer. *CMRO* 2006;22(10):1949-54.
35. Dagtekin O, Berlet T, Delis A, **Kampe S**. Manually controlled total intravenous anesthesia augmented by electrophysiologic monitoring for complex stereotactic neurosurgical procedures. *J Neurosurg Anesthesiol* 2007;19(1):45-8.
36. Picker SM, Kaleta T, Hekmat K, **Kampe S**, Gathof B. Antiplatelet therapy preceding coronary artery surgery: Implications for bleeding, transfusion requirements, and outcome. *Eur J Anaesthesiol* 2007;24(4):332-9.
37. Dagtekin O, Hotz A , **Kampe S**, Auweiler M, Warm M. Postoperative analgesia and flap perfusion after pedicled TRAM flap reconstruction - continuous wound instillation with ropivacaine 0.2%. *Plastic and Reconstructive Surgery* 2008 (in press).